

# **THE MASS OF THE CLASSICAL CEPHEID S MUS**

**NASA Grant No. NAG5-11946**

**Final Report**

**For Period 1 April 2002 through 31 March 2004**

**Principal Investigator  
Dr. Nancy Evans**

**August 2004**

**Prepared for:**

**National Aeronautics and Space Administration  
Goddard Space Flight Center  
Greenbelt, MD 20771**

**Smithsonian Institution  
Astrophysical Observatory  
Cambridge, Massachusetts 02138**

<p><b>The Smithsonian Astrophysical Observatory is a member of the Harvard-Smithsonian Center for Astrophysics</b></p>
--

**The NASA Technical Officer for this Grant is George Sonneborn, Code 681,  
NASA/Goddard Space Flight Center, Greenbelt, MD 20771**

A FUSE spectrum of the hot companion of the Cepheid S Mus has been obtained, reduced and analysed. Since the determination of the temperature of S Mus B is the prime goal of this program, we have identified two regions in the spectrum which are temperature sensitive in the temperature range of the companion, but which are not overly affected by molecular hydrogen absorption. From comparison of the spectrum with spectra of stars of known temperature, we have determined the temperature of the companion to be 17,000 K. From this a mass has been inferred. Combining this mass with the orbital velocity ratio of the Cepheid and the companion (from the ground-based Cepheid orbit, and HST GHRS echelle observations), the mass of the Cepheid is determined to be  $6.0 \pm 0.4$  solar masses. The results have been presented at a conference (conference proceedings in press--see attachment). Final paper is nearing journal submission.

## Cepheid Masses: FUSE Observations of S Mus

Nancy Ramage Evans<sup>1</sup>, Derck Massa<sup>2</sup>, Alexander Fullerton<sup>3</sup>,  
George Sonneborn<sup>4</sup>, Rosina Iping<sup>4</sup>

<sup>1</sup>*Smithsonian Astrophysical Observatory, 60 Garden St., Cambridge,  
MA 02138, USA*

<sup>2</sup>*NASA's GSFC, SGT, Inc., Code 681.0, Greenbelt, MD 20771, USA*

<sup>3</sup>*University of Victoria, Department of Physics and Astronomy, PO Box  
3055, Station CSC, Victoria, BC, Canada*

<sup>4</sup>*NASA's GSFC, Code 681, Greenbelt, MD 20771, USA*

**Abstract.** S Mus is the Cepheid with the hottest known companion. As a benefit, the large ultraviolet flux made it the only Cepheid companion for which the velocity amplitude could be measured with the echelle mode of the HST GHRS. Unfortunately, the high temperature is difficult to constrain at wavelengths longer than 1200 Å because of the degeneracy between temperature and reddening. We have now obtained a FUSE spectrum in order to determine the temperature of the companion better. We have identified two regions which are temperature sensitive near 16 000 K, but are relatively unaffected by H<sub>2</sub> absorption (940 Å, and the Ly β wings). By comparing S Mus B with other FUSE spectra, we have determined a temperature of 17 000 K. The resultant Cepheid mass is  $6.0 \pm 0.4 M_{\odot}$ .

### 1. Introduction

Measurement of Cepheid masses has been a goal both for a thorough understanding of these primary distance indicators and also as a benchmark for stellar evolution calculations of high mass stars. Specifically, the most important uncertainty in the calculations of evolutionary tracks of massive stars near the main sequence, core convective overshoot, can be constrained by the combination of a measured Cepheid mass and its luminosity.

Ultraviolet high resolution spectroscopy has provided a group of Cepheid double-lined spectroscopic binaries. Specifically, the orbital velocity amplitudes of hot companions can be measured with the Hubble Space Telescope (HST) Space Telescope Imaging Spectrograph (STIS) or Goddard High Resolution Spectrograph (GHRS) and formerly IUE. The ratio of this amplitude with the orbital velocity amplitude of the Cepheid from a ground-based orbit provides the inverse mass ratio. Typically, a very accurate temperature or spectral type for the hot companion can be obtained from IUE low resolution spectra from

1200 to 3200 Å, from which the mass of the companion can be inferred very accurately.

The first step in the process is to determine the orbits of the Cepheids from the ground. As a sidelight, from a recent compilation of Cepheid orbits (Evans et al. 2004) approximately half the Cepheid binaries have been found to be triple systems. There is some bias in this fraction, since a massive secondary will increase the probability that the binary motion of the Cepheid is detected and studied. However, Cepheids have proved a valuable source of information about the multiplicity of massive stars.

The Cepheid S Mus, however, has required a different approach than other Cepheid binaries. For a summary, see Evans et al. (1998, 2004). Its companion is the hottest known in a Cepheid system. This allowed an observation with the HST GHRS echelle mode (Böhm-Vitense et al. 1997), providing the most accurate velocity available for a binary Cepheid. However, this high temperature made it difficult to determine the temperature of the companion. Specifically, the energy distribution of the spectrum rises monotonically in the IUE region (1200 to 3200 Å). This means the changes in the spectrum with temperature are small, and also that the temperature is coupled with reddening. The first step in remedying this was to obtain a Voyager spectrum of S Mus B (Evans, Holberg, & Polidan 1996). The Voyager spectral region (to wavelengths as short as 900 Å) is complicated by the H<sub>2</sub> absorption. Because Voyager spectra have low resolution, only approximate corrections can be applied to correct for the H<sub>2</sub> absorption.

We recently obtained a FUSE spectrum in the 900 to 1200 Å region. The aim of the project was to identify features which are temperature sensitive in the relevant spectral range (B3V to B5V) which are only slightly affected by H<sub>2</sub> lines. We have used spectra from the FUSE archive (Table 1) to look for appropriate features. The spectra were *reddened* to the  $E(B - V)$  of S Mus ( $E(B - V) = 0.23$  which corresponds to a column density of  $N_H = 1.4 \times 10^{21} \text{ cm}^{-2}$ ) and then normalized using the flux in the wavelength range 1140 to 1170 Å. The comparison stars are listed in Table 1, together with the temperatures and gravities determined by one of us (DM) from Strömgren photometry, using the calibration of Napiwotzki et al. (1993).

Table 1. Comparison Stars

Star	Spectral Type	$E(B - V)$ (mag)	T (K)	$\log g$ $\text{cm s}^{-2}$
HD 97991	B1 V	0.02	25 500	4.00
HD 121800	B1.5 V	0.07	20 300	1.30
HDE 233622	B2 V	0.03	21 000	3.60
HD 51013	B3 V	0.00	17 100	4.30
HD 261878	B3 V	0.04	15 500	4.50
HD 72350	B4 IV	0.16	15 650	3.00
HD 35899	B5 V	0.03	16 700	4.30
HD 37332	B5 V	0.02	15 600	4.50
HD 37641	B6 V	0.05	12 800	4.40




Figure 1. Ly  $\beta$  for the Cepheid companion S Mus B (solid line) compared with HD 51013 ( $T = 17\,100$  K; dotted line) and HD 37332 ( $T = 15\,600$  K; dashed line). The dot-dash line is the H<sub>2</sub> absorption, showing that outside the line core; the wings are relatively unaffected by H<sub>2</sub> absorption.

Two regions were identified for temperature determination:  $\text{Ly}\beta$  and the region near 940 Å, the shortest wavelength where flux is apparent in the S Mus B spectrum. Fig. 1 shows the comparison between the  $\text{Ly}\beta$  for S Mus B and the comparison stars, HD 51013 (B3V) and HD 37332 (B5V).  $\text{Ly}\beta$  is gravity sensitive, but we have restricted our comparisons to main sequence stars to remove that dependency. S Mus B is clearly a very good match to HD 51013 (B3V), and markedly different from HD 37332 (B5V). The same is true in the 940 Å region. Based on this, we adopt 17 000 K as the temperature, with an uncertainty of  $\pm 500$  K.

In order to derive a mass corresponding to this temperature we have used the masses from Andersen (1991). We have combined these with the more recent temperatures from Ribas et al. (2000), based on Strömgren photometry, which should be comparable to the temperatures for our standard stars. The mass for S Mus B from the FUSE temperature and a linear fit to early B stars from the Andersen Ribas set is  $5.3 \pm 0.26 M_{\odot}$ .

The resulting mass from the previously measured velocity amplitude ratio ( $1.14 \pm 0.06$ , Böhm-Vitense et al. 1997) is  $6.0 \pm 0.4 M_{\odot}$ . This is very similar to the previous mass for the Cepheid (Böhm-Vitense et al. 1997), but with a significantly smaller uncertainty.

## References

- Andersen, J. 1991, ARA&A, 3, 91  
 Böhm-Vitense, E., Evans, N.R., Carpenter, K., Beck-Winchatz, B., Robinson, R. 1997, ApJ, 477, 916  
 Evans, N.R., Böhm-Vitense, E., Carpenter, K., Beck-Winchatz, B., Robinson, R. 1998, ApJ, 494, 768  
 Evans, N.R., Holberg, J.B., Polidan, R.S. 1996, AJ, 112, 264  
 Evans, N.R., Carpenter, K., Robinson, R., Dekas, A., Kienzle, F. 2004, in preparation  
 Napiwotzki, R., Schönberner, D., Wenske, V. 1993, A&A, 268, 653  
 Ribas, I. 2000, MNRAS, 313, 99

## Discussion

*Aerts:* How large is your convective overshooting parameter that fits best for S Mus?

*Evans:* The mass of S Mus is close to that predicted by the moderate overshoot (0.2 pressure scale heights) of both the Geneva group (Schaller et al. 1992, A&AS, 96, 269) and the Padua group (Bertelli et al. 1994, A&AS, 106, 275).

*Kawaler:* Could you clarify what core overshoot is being constrained? Cepheids have He burning cores which are convective.

*Evans:* It is the boundary during the main sequence phase between the convective core and the radiative envelope. This determines the amount of fuel available during H burning, the main sequence lifetime, and also the luminosity during the He burning blue loops.